

## Minimal Invasive Dentistry: Foundational Articles and Consensus Recommendations, 2020

**Crystal YO, Marghalani AA, Ureles SD, Wright JT, et al.** Use of silver diamine fluoride for dental caries management in children and adolescents, including those with special health care needs. Pediatr Dent. 2017, 15; 39(5): 135-145

**Chatzimarkou S, Koletsi D, Kavvadia K.** The effect of resin infiltration on proximal caries lesions in primary and permanent teeth. A systematic review and meta-analysis of clinical trials. Journal Dentistry 2018; 77: 8-17.

**Damian H, Monika HZ, Spyridon NP, Theodore E.** Interventions for orthodontically induced white spot lesions: a systematic review and meta-analysis. Eur J Orthodont, 2017; 39(2):122-133.

Feartherstone JBD. Dental caries – a dynamic disease process. Australian Dent J, 2008; 53(3): 286-291.

**Frencken JE, Peters Mc, Manton DJ, et al.** Minimal intervention dentistry for managing dental caries. Int Dent J. 2012 62(5): 223-243.

Innes NPT, ChuCH, Fontana M, Lo ECM, Thomson WM, Uribe S, Heiland M, Jepsen S, Schwendicke F. A century of change towards prevention and minimal intervention in cariology. J Dent Res, 2019; 98(6): 611-617.

**Innes NPT, Frencken JE, Bjørndal L et al.** Managing carious lesions: consensus recommendations on terminology. Adv Dent Research 2016; 28(2): 49-57.

**Jingawar MM, Bajwa NK, Palhak A.** Minimal intervention dentistry – a new frontier in clinical dentistry. J Clin Dent Res, 2014; 8(7): ZE04- ZE08.

Li T, Zhai X, Song F, Zhu H. Selective versus non-selective removal for dental caries: a systematic review and meta-analysis. Acta Odontol Scand, 2018; 76 (2): 135-140.

Liang Y, Deng Z, Dai 1 X, Tian J, Zhao W. Micro-invasive interventions for managing non-cavitated proximal caries of different depths: a systematic review and meta-analysis. Clin Oral Investig, 2018; 22: 2675–2684.

Schwendicke F, Frencken JE, Bjorndal et al. Managing carious lesions. Adv Dent Research, 2016; 28(2): 49-57.

**Coelho Leal, S.** Minimal intervention dentistry in the management of the paediatric patient. Brit Dent J 2014; 216(11):623-7.

**Trieu A, Mohamed A, Lynch E.** Silver diamine fluoride versus sodium fluoride for arresting dentine caries in children: a systematic review and meta-analysis. Scientific Reports, 2018; 9: 115-224.

**Tyas MJ, Anusavice KJ, Frencken JE, Mount GJ.** Minimal intervention dentistry – a review. FDI Commission Project 1-97. Int Dent J 2000; 50: 1-12.

Weerheijm KL, Groen HJ. The residual caries dilemma. Community Dent Oral Epidemiol, 1999; 27: 436-441.

## IAPD Consensus Recommendations

**1.** Minimal invasive dentistry focuses on caries arrest procedures rather than surgical intervention.

**2.** The strategies for minimal invasive dentistry include: early caries detection and caries risk assessment; remineralization of demineralized enamel and dentin; optimal caries preventive measures; minimally invasive surgical interventions; and repair rather than replacement of restorations.

**3.** Demineralized, non-cavitated carious lesions may naturally arrest or arrest through preventive interventions.

**4.** The use of silver diamine fluoride (SDF) can be effective in arresting cavitated caries lesions.

**5.** Micro-invasive approaches such as infiltration methods, can be effective in the management of non-cavitated proximal carious lesions.

6. The use of minimal invasive dentistry should be complemented by preventive measures for caries arrest that include optimization of patient's topical fluoride exposure, and maintenance of good oral health practices.

**7.** When necessary, surgical interventions should emphasize minimal cavity designs, conservative removal of carious dentin approximating the pulp, and adhesive restorative materials.

How to cite: IAPD Foundational Articles and Consensus Recommendations: Minimal Invasive Dentistry, 2020. http://www.iapdworld.org/06\_minimal-invasive-dentistry.